

Patient information leaflet

Sublingual Immunotherapy (SLIT) for House Dust Mite with ACARIZAX: General Information

What is sublingual immunotherapy?

There are two ways of re-training the immune system to tolerate a substance the individual is allergic to (desensitisation): by either administering allergen extract under the skin (subcutaneous immunotherapy, SCIT) or under the tongue (sublingual immunotherapy, SLIT). The sublingual (under the tongue) route is convenient (as it does not involve injections) and very safe, provided the instructions about product administration are followed correctly.

What is ACARIZAX?

ACARIZAX is a sublingual (under the tongue) immunotherapy treatment suitable for patients with severe allergic rhinitis symptoms caused by House Dust Mite allergy. ACARIZAX is a dispersible tablet that contains a measured amount of 4 different House Dust Mite proteins.

Can anybody take ACARIZAX?

ACARIZAX cannot be given to children younger than 5 years and to individuals with certain medical conditions or who are on certain medications (severe asthma, fish allergy, beta blocker medications, to name a few). Your allergist will make an individual assessment and decide if ACARIZAX can be considered as a treatment option. ACARIZAX is unlikely to benefit individuals with milder allergic rhinitis symptoms and should only be considered if symptoms are severe and the usual allergic rhinitis medications are not working despite regular use.

What are the benefits of taking ACARIZAX?

The treatment aims to re-train the immune system of the allergic individual to tolerate House Dust Mite better. At the end of the treatment period the patient can expect around reduction in symptoms caused by House Dust Mite allergy and reduction in the regular medications taken to relieve these symptoms (e.g. antihistamines, nasal sprays etc). The treatment is not a complete cure, as it does not hold the promise to eliminate the House Dust Mite allergy altogether, but helps tolerise the individual to a point where the allergy symptoms are much milder. The benefit of the treatment is sustained for a number of years after cessation of immunotherapy. In individuals who do not already have asthma, immunotherapy can also reduce the risk of future asthma.

How is ACARIZAX taken?

The very first dose is given in hospital, to ensure it is well tolerated. During this visit you will be given instructions about how to continue administering the medication at home.

The tablet should be taken every day at a fixed time. The tablet is simply placed under the tongue with clean dry hands. It melts almost immediately, but the patient is instructed not to swallow for 1-2 min so that the extract is absorbed via the lining of the mouth. After 1-2 minutes the patient can swallow as normal.

Can ACARIZAX treatment be started at any time?

ACARIZAX can be started at any time of the year, provided the patient is otherwise well.

How long is ACARIZAX taken for?

To get the maximum benefit from this treatment, ACARIZAX should be taken every day for 3 years. The dose remains the same for the whole duration of treatment (1 tablet a day).

There are certain conditions, such as feverish illness and dental extractions, where the treatment should be discontinued for a short time until the patient is recovered, after which the treatment should be re-started promptly. You/your child will be instructed about these exceptional conditions and when and how to re-start treatment.

What are the possible side effects of ACARIZAX?

The most common side effect of the treatment is tingling or itching sensation under the tongue, in the mouth, throat or ears. This tends to happen immediately after the tablet is administered and usually lasts for no more than 10-15 min. Itch and tingling can be managed by taking an

antihistamine 30min before taking the ACARIZAX tablet. These complaints usually subside once the body gets used to the treatment, within 2-3 weeks. However, if they persist you should discuss this with your allergist.

Other rarer side effects include headaches, tiredness, swelling of the tongue or lips, itchy hives on skin, eczema flare, sneezing, runny or blocked nose and watery itchy eyes. You should take antihistamine if these complaints occur and inform your allergist.

Very rarely the patient may experience more severe symptoms such as difficulty swallowing, hoarse voice, feeling of swollen throat, difficulty breathing, wheeze or feeling unwell and dizzy. These symptoms would be exceptionally unlikely to occur if the medication has been tolerated well initially and is taken in the correct way.

How long will the beneficial effect of ACARIZAX last?

This varies from one individual to another. In general, the effect of this treatment is expected to last for several years after the 3 year course of treatment is completed.

Should I stop taking my other hay fever medications if I am on ACARIZAX?

No. For the first 6 months after you have started ACARIZAX you must remain on the same regular allergic rhinitis medications you were taking before. Your doctor will guide you about how to reduce your regular allergic rhinitis medications at follow up in clinic visits, depending on the severity of your symptoms.

Patient information leaflet

Sublingual Immunotherapy (SLIT) for House Dust Mite with ACARIZAX -Specific information about treatment

How to take ACARIZAX

- NEVER take more than one tablet per day.
- Do not exercise for 1 hour before administration.
- Take the medication at a fixed time every day, ideally in the morning.
- Ensure patient's mouth is clean of food, rinse mouth if necessary.
- Remind/remember the need to hold the tablet under the tongue for 1-2 min before swallowing.
- Ensure hands are clean and DRY.
- Tear off the strip marked with triangles at the top of the pack.
- Tear a square off the pack along the perforated lines.
- DO NOT force the tablet through the foil. It may damage the tablet as it easily breaks. Instead, fold back the marked corner of the foil and then pull it off.
- Remove the tablet carefully from the foil with dry hands.
- Take tablet immediately by opening the mouth, lifting tongue up and gently placing the tablet under the tongue.
- Do not swallow for 1-2 min to allow the extract to be absorbed through the lining of the mouth under the tongue.
- Do not eat or drink anything for the next 5-10 minutes after administration.
- Do not exercise for 1 hour after administration.
- The extract in the tablet needs to be absorbed to the body through the lining of the mouth under the tongue. Therefore, NEVER chew the tablet and NEVER swallow the tablet directly.

When should ACARIZAX be stopped?

ACARIZAX should be stopped if any of the following happens:

1. Asthma attack
2. Infection
3. Fever
4. On the day another vaccination is given and whilst experiencing the side effects of another vaccination

ACARIZAX should be restarted once the patient has recovered completely from the above listed conditions.

5. Infections/inflammation/bleeding/wounds/ulcers in the mouth (for example, cold sores, dental extraction, trauma to mouth during sports injury)
ACARIZAX should be restarted after the wounds in the mouth have healed and 7 days after tooth extraction.

How should ACARIZAX be re-started after a break or a missed dose?

DO NOT take a double dose.

If the treatment is interrupted for less than 10-14 days the treatment can be re-started at home. Simply take your 1 tablet per day as usual.

If the treatment has been interrupted for more than 10-14 days then treatment should be re-started under supervision in the hospital. Please inform your allergist or allergy specialist nurse.

How to store ACARIZAX

Keep out of reach of children.

Store in safe dry place at room temperature. You do not need to keep ACARIZAX in the refrigerator.

Open each blister just before you take the tablet. DO NOT take tablets out of the blisters to store in a pillbox.

Repeat prescription

It is the family's responsibility to phone the provider of repeat ACARIZAX prescriptions well in advance before running out of tablets. In order to avoid interruption of treatment we advise that the family phones to re-order next prescription on the day they open the last 30 tablet pack.

Could any new medications be started whilst on ACARIZAX?

You MUST discuss with your allergist before starting any new medications or having other vaccines. **At any time during the treatment period of 3 years, PLEASE REMEMBER to inform your allergist of any new medications you are advised to take.**

YOUR ALLERGY TEAM CONTACTS ARE:

Dr Sibel Sonmez-Ajtai

Consultant in Allergy

Secretary Tel: (0114) 271 7585

Mobile: 07557564254

E-mail: sibel.sonmez-ajtai@nhs.net

Dr Nicola Jay

Consultant in Allergy

and Asthma

Secretary Tel: (0114) 271 7585

E-mail: nicola.jay@nhs.net

Dr Eleanor Minshall

Consultant in Allergy

Secretary Tel: (0114) 271 7585

E-mail: eleanor.minshall@nhs.net

Allergy Specialist Nurses

Tel: (0114) 226 7872